

# Lie symmetry analysis and conservation laws for the time fractional fourth-order evolution equation

Wang Li, Tian Shoufu\*, Feng Lianli, Song Xiaoqiu\*

(School of Mathematics, China University of Mining and Technology, Xuzhou 221116, China)

**Abstract:** In this paper, we study Lie symmetry analysis and conservation laws for the time fractional nonlinear fourth-order evolution equation. Using the method of Lie point symmetry, we provide the associated vector fields, and derive the similarity reductions of the equation, respectively. The method can be applied wisely and efficiently to get the reduced fractional ordinary differential equations based on the similarity reductions. Finally, by using the nonlinear self-adjointness method and Riemann-Liouville time-fractional derivative operator as well as Euler-Lagrange operator, the conservation laws of the equation are obtained.

**Key words:** Lie symmetry method; symmetry analysis; conservation laws

**CLC number:** O 175.12      **Document code:** A      **Article ID:** 1000-5137(2017)03-0363-12

**2000 MSC:** 35Q51, 35Q53, 35C99, 68W30, 74J35

## 四阶时间分数阶演化方程的 Lie 对称分析和守恒律

王 丽, 田守富\*, 冯连莉, 宋晓秋\*

(中国矿业大学 数学学院, 徐州 221116)

**摘 要:** 主要研究了四阶时间分数阶演化方程的 Lie 对称分析和守恒. 基于 Lie 点对称方法, 分别得到了该方程的相关向量场以及相似约化. 在相似约化的基础上, 通过该方法来获得分数阶常微分方程是非常有效的. 最后, 通过非线性的自伴随方法和时间分数阶的黎曼-刘维尔导数算子以及欧拉-拉格朗日算子, 得到了该方程的守恒律.

**关键词:** Lie 对称方法; 对称分析; 守恒律

---

**Received date:** 2016-09-12

**Foundation item:** This research was supported by the National Training Programs of Innovation and Entrepreneurship for Undergraduates (201410290039); the Fundamental Research Funds for the Central Universities (2015QNA53, 2015XKQY14); the General Financial Grant from the China Postdoctoral Science Foundations (2015M570498 and 2017T100413); Natural Sciences Foundation of China (11301527).

\***Corresponding author:** Tian Shoufu, associate professor, reseach area: partial differential equation. E-mail: sf-tian@cumt.edu.cn, shoufu 2006@126.com; Song Xiaoqiu, professor, reseach area: functional analysis. E-mail: songxiaoqiu@cumt.edu.cn

## 1 Introduction

As we all know, the Lie group of transformations theory, was initially constructed by the Norwegian mathematician Sophus Lie in the early 19th century, which has been widely studied in differential equations<sup>[1–12]</sup>, particularly in partial differential equations (PDEs) with multi-component systems. In recent years, it is important to investigate the fractional differential equations (FDEs) because of more and more applications in various fields<sup>[13–16]</sup>: systems identification, fluid flow, control problems, signal processing, viscoelastic materials, polymers, fluid mechanics, physics, economics, biology, engineering and so on. Actually, the symmetry analysis of FDEs and the fractional derivatives are proposed by Gazizov and his collaborators in [17]. Readers can refer to [18–22] for further details. In [23], the famous Noether theorem connects the symmetry theory with conservation laws of differential equations, which exist Euler-Lagrange equations. In addition, in order to look for conservation laws of FDEs, Frederico et al.<sup>[24–28]</sup> have provided fractional generalizations of Noether theorem. However, it's limited for many FDEs to deal with fractional generalizations. Then, a general theorem on conservation laws for arbitrary differential equations<sup>[29]</sup> was proposed by Ibragimov. Based on it, Lukashchuk<sup>[30]</sup> proposed Noether operators without fractional Lagrangians and proved fractional generalizations of the Noether operator. In [31], in order to get the conservation laws, the authors applied the fractional generalizations of the Noether operators to the time-fractional Kompaneets equations.

In this paper, we study the Lie symmetry analysis to the time fractional nonlinear fourth-order evolution equation

$$\frac{\partial^\alpha u}{\partial t^\alpha} + h_1 u_x u_{xx} + h_2 u^2 u_{xx} + h_3 u u_x^2 + h_4 u_{xxxx} = 0, \quad 0 < \alpha \leq 2, \quad (1)$$

where  $u = u(x, t)$ ,  $h_1, h_2, h_3$  and  $h_4$  are constants. When  $\alpha = 1$ ,  $h_1 = -2$ ,  $h_2 = -1$ ,  $h_3 = -2$ ,  $h_4 = 1$ , the time fractional equation becomes the following nonlinear fourth-order evolution equation

$$u_t - 2u_x u_{xx} - u^2 u_{xx} - 2u u_x^2 + u_{xxxx} = 0. \quad (2)$$

Some new exact solutions of Eq. (2) are given in [32].

The paper is divided into five parts. In Section 2, some properties analyzing FDEs of Lie group method, and the associated infinitesimal generator are introduced. In Section 3, we obtain the corresponding similarity reduction of Eq. (1). Section 4 is devoted to construct the conservation laws. Finally, some conclusions and discussions are present.

## 2 Lie symmetry analysis

In this section, we investigate the time fractional nonlinear fourth-order evolution equation by means of Lie symmetry. First we recall, some basic properties of Lie symmetry analysis of FDEs. Here is the definition of Riemann-Liouville fractional partial derivative in [21].

**Definition 1**<sup>[21]</sup> The Riemann-Liouville fractional partial derivative is defined by

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \begin{cases} \frac{\partial^m u}{\partial t^m}, & (\alpha = m \in \mathbb{N}), \\ \frac{1}{\Gamma(m - \alpha)} \frac{\partial^m}{\partial t^m} \int_0^t \frac{u(\tau, x)}{(t - \tau)^{\alpha + 1 - m}} d\tau, & (m - 1 < \alpha < m, \quad m \in \mathbb{N}), \end{cases} \quad (3)$$

where  $\partial_t^m$  is the usual partial derivative of integer order  $m$ .

Consider a time fractional FDE with two independent variables

$$\frac{\partial^\alpha u(x, t)}{\partial t^\alpha} = F(x, t, u, u_x, u_{xx}, \dots), \alpha > 0. \tag{4}$$

One-parameter point transformations of Lie group is given by

$$\begin{aligned} x^* &= x + \epsilon \xi(x, t, u) + O(\epsilon^2), \\ t^* &= t + \epsilon \tau(x, t, u) + O(\epsilon^2), \\ u^* &= u + \epsilon \eta(x, t, u) + O(\epsilon^2), \\ \frac{\partial^\alpha u^*}{\partial t^{*\alpha}} &= \frac{\partial^\alpha u}{\partial t^\alpha} + \epsilon \eta^{\alpha t}(x, t, u) + O(\epsilon^2), \\ \frac{\partial^j u^*}{\partial x^{*j}} &= \frac{\partial^j u}{\partial x^j} + \epsilon \eta^{jx}(x, t, u) + O(\epsilon^2), j = 1, 2, \dots, \end{aligned} \tag{5}$$

where  $\epsilon$  is the group parameter, and  $\xi, \tau, \eta$  are infinitesimals. In addition, we consider the following infinitesimal generator

$$V = \tau(x, t, u) \frac{\partial}{\partial t} + \xi(x, t, u) \frac{\partial}{\partial x} + \eta(x, t, u) \frac{\partial}{\partial u}, \tag{6}$$

here,

$$\tau(x, t, u) = \left. \frac{dt^*}{d\epsilon} \right|_{\epsilon=0}, \quad \xi(x, t, u) = \left. \frac{dx^*}{d\epsilon} \right|_{\epsilon=0}, \quad \eta(x, t, u) = \left. \frac{du^*}{d\epsilon} \right|_{\epsilon=0}. \tag{7}$$

Based on the infinitesimal invariance criterion, we find the infinitesimal generator must satisfy

$$pr^{(\alpha,4)}V(\Delta_1)|_{\Delta_1=0} = 0, \tag{8}$$

where  $\Delta_1 = u_t^\alpha + h_1 u_x u_{xx} + h_2 u^2 u_{xx} + h_3 u u_x^2 + h_4 u_{xxx}$ .

The prolongation operator  $pr^{(\alpha,4)}$  is

$$pr^{\alpha,4}V = V + \xi_\alpha^0 \partial_{t^\alpha} u + \eta^x \partial_{u_x} + \eta^{xx} \partial_{u_{xx}} + \eta^{xxx} \partial_{u_{xxx}} + \eta^{xxxx} \partial_{u_{xxxx}}, \tag{9}$$

here,

$$\begin{aligned} \eta^x &= \eta_x + (\eta_u - \xi_x)u_x - \tau_x u_t - \xi_u u_x^2 - \tau_u u_x u_t, \\ \eta^{xx} &= \eta_{xx} + (2\eta_{xu} - \xi_{xx})u_x - \tau_{xx} u_t + (\eta_{uu} - 2\xi_{xu})u_x^2 - 2\tau_{xu} u_x u_t - \xi_{uu} u_x^3 \\ &\quad - \tau_{uu} u_x^2 u_t + (\eta_u - 2\xi_x)u_{xx} - 2\tau_x u_{xt} - 3\xi_u u_{xx} u_x - \tau_u u_{xx} u_t - 2\tau_u u_{xt} u_x, \\ \eta^{xxx} &= \eta_{xxx} + (3\eta_{xuu} - \xi_{xxx})u_x - \tau_{xxx} u_t + 3(\eta_{xuu} - \xi_{xxu})u_x^2 - 3\tau_{xuu} u_x u_t \\ &\quad + (\eta_{uuu} - 3\xi_{xuu})u_x^3 + 3(\eta_{xu} - \xi_{xx})u_{xx} - 3\tau_{xx} u_{xt} - 3\tau_{xuu} u_x^2 u_t + 3(\eta_{uu} - 3\xi_{xu})u_x u_{xx} \\ &\quad - 3\tau_{xu} u_{xx} u_t - 6\tau_{xu} u_{xt} u_x - 3\tau_x u_{xxt} + (\eta_u - 3\xi_x)u_{xxx} - \xi_{uuu} u_x^4 - 6\xi_{uu} u_{xx} u_x^2 - 3\tau_{uu} u_x^2 u_{xt} \\ &\quad - \tau_{uuu} u_x^3 u_t - 3\xi_u u_{xx}^2 - 3\tau_u u_{xxt} u_x - 3\tau_u u_{xt} u_{xx} - 3\tau_{uu} u_t u_x u_{xx} - 4\xi_u u_x u_{xxx} - \tau_u u_t u_{xxx}, \\ \eta^{xxxx} &= \eta_{xxxx} + u_x(4\eta_{xxuu} - \xi_{xxxx}) - \tau_{xxxx} u_t + u_x^2(6\eta_{xxuu} - 4\xi_{xxxu}) + u_{xx}(6\eta_{xuu} - 4\xi_{xxx}) - 4\tau_{xxuu} u_t u_x \\ &\quad - 4\tau_{xxx} u_{xt} + u_x^3(4\eta_{xuuu} - 6\xi_{xuuu}) + u_x u_{xx}(12\eta_{xuu} - 18\xi_{xuu}) - 6\tau_{xuu} u_x^2 u_t - 6\tau_{xuu} u_{xx} u_t \\ &\quad - 12\tau_{xuu} u_x u_{xt} + u_x^4(\eta_{uuuu} - 4\xi_{uuuu}) + u_x^2 u_{xx}(6\eta_{uuu} - 24\xi_{uuu}) + u_{xxx}(4\eta_{xu} - 6\xi_{xx}) - 6\tau_{xx} u_{xxt} \end{aligned}$$

$$\begin{aligned}
& -4\tau_{xuuu}u_x^3u_t - 12\tau_{xuu}u_tu_xu_{xx} - 12\tau_{xuu}u_x^2u_{xt} + (3\eta_{uu} - 12\xi_{xu})u_{xx}^2 + u_xu_{xxx}(4\eta_{uu} - 16\xi_{xu}) \\
& - 12\tau_{xu}u_{xx}u_{xt} - 4\tau_{xu}u_tu_{xxx} - 12\tau_{xu}u_xu_{xxt} - 4\tau_xu_{xxx} + u_{xxx}(\eta_u - 4\xi_x) - \xi_{uuuu}u_x^5 \\
& - 10\xi_{uuu}u_x^3u_{xx} - 15\xi_{uu}u_xu_{xx}^2 - 10\xi_{uu}u_x^2u_{xxx} - 4\tau_{uuu}u_x^3u_{xt} - 12\tau_{uu}u_xu_{xx}u_{xt} - 6\tau_{uu}u_x^2u_{xxt} \\
& - \tau_{uuuu}u_x^4u_t - 6\tau_{uuu}u_x^2u_{xx}u_t - 10\xi_uu_{xxx} - 6\tau_uu_{xxt} - 4\tau_uu_{xxt} \\
& - 4\tau_uu_{xt}u_{xxx} - 3\tau_{uu}u_tu_{xx}^2 - 4\tau_{uu}u_tu_xu_{xxx} - 5\xi_uu_xu_{xxx} - \tau_uu_tu_{xxx}.
\end{aligned} \tag{10}$$

From [18–21], we have

$$\begin{aligned}
\xi_\alpha^0 &= \frac{\partial^\alpha \eta}{\partial t^\alpha} + (\eta_u - \alpha D_t(\tau)) \frac{\partial^\alpha u}{\partial t^\alpha} - u \frac{\partial^\alpha \eta_u}{\partial t^\alpha} \\
&+ \mu + \sum_{n=1}^{\infty} \left[ \binom{a}{n} \frac{\partial^\alpha \eta_u}{\partial t^\alpha} - \binom{a}{n+1} D_t^{n+1}(\tau) \right] \\
&\times D_t^{\alpha-n}(u) - \sum_{n=1}^{\infty} \binom{a}{n} D_t^n(\xi) D_t^{\alpha-n}(u_x),
\end{aligned} \tag{11}$$

where

$$\mu = \sum_{n=2}^{\infty} \sum_{m=2}^n \sum_{k=2}^m \sum_{r=0}^{k-1} \binom{a}{n} \binom{n}{m} \binom{k}{r} \frac{1}{k!} \frac{t^{n-\alpha}}{\Gamma(n+1-\alpha)} [-u]^r \frac{\partial^m}{\partial t^m} [u^{k-r}] \frac{\partial^{n-m+k} \eta}{\partial t^{n-m} \partial u^k}, \tag{12}$$

with the additional constraint condition

$$\tau(x, t, u)|_{t=0} = 0. \tag{13}$$

In addition, when the infinitesimal  $\eta$  is linear to  $u$ ,  $\mu$  is invalid [19].

Now, we apply the above Lie symmetry analysis method above all to the Eq.(1), and obtain the following theorem.

**Theorem 1** The vector fields of the time fractional fourth-order evolution equation is given by

$$V_1 = x \frac{\partial}{\partial x} + \frac{4t}{\alpha} \frac{\partial}{\partial t} - u \frac{\partial}{\partial u}, \quad V_2 = \frac{\partial}{\partial x}. \tag{14}$$

**Proof** Substituting (9) and (10) into (1), we find the following symmetry equation

$$\xi_\alpha^0 + h_4 \eta^{xxxx} + h_1 \eta^{xx} u_x + h_2 u^2 \eta^{xx} + h_1 \eta^x u_{xx} + 2h_3 u \eta^x u_x + 2h_2 u \eta u_{xx} + h_3 \eta u_x^2 = 0, \tag{15}$$

by the condition that variables  $u_x, u_t, u_{xt}, u_{xx}, u_{tt}, u_{xtt}, \dots$  and  $D_t^{\alpha-n}u, D_t^{\alpha-n}u_x$  for  $n = 1, 2, \dots$  of  $u$  are independent. Taking (10)-(13) into consideration, and making each power of the derivatives of  $u$  is equal to 0, we get the determining equations

$$\begin{aligned}
& \xi_u = \xi_t = \tau_u = \tau_x = \eta_{uu} = 0, \\
& h_4(\alpha\tau_t - 4\xi_x) = 0, \\
& \frac{\partial^\alpha \eta}{\partial t^\alpha} - u \frac{\partial^\alpha \eta_u}{\partial t^\alpha} + h_2 \eta_{xx} u^2 = 0, \\
& h_4(12\eta_{xuu} - 18\xi_{xuu}) + h_1(\eta_u + \alpha\tau_t - 3\xi_x) - 3h_2 \xi_u u^2 = 0, \\
& \binom{a}{n} \frac{\partial^\alpha \eta_u}{\partial t^\alpha} - \binom{a}{n+1} D_t^{n+1}(\tau) = 0, \text{ for } n = 1, 2, \dots
\end{aligned} \tag{16}$$

Solving (16), we find

$$\begin{aligned}\xi &= a_1x + a_2, \\ \tau &= \frac{4a_1}{\alpha}t, \\ \eta &= -a_1u,\end{aligned}\tag{17}$$

where,  $a_1, a_2$  are arbitrary constants. So the associated vector fields are given by

$$V_1 = x \frac{\partial}{\partial x} + \frac{4t}{\alpha} \frac{\partial}{\partial t} - u \frac{\partial}{\partial u}, \quad V_2 = \frac{\partial}{\partial x}.\tag{18}$$

### 3 The similarity reduction

In this section, the similarity reduction for the time fractional nonlinear fourth-order evolution equation is obtained. Similar to the definition of group invariant solution of PDEs<sup>[4, 33]</sup>, the group invariant solution of FDEs is defined.

**Definition 2**<sup>[4, 33]</sup> If  $u = v(x, t)$  satisfies Eq. (1), and  $u = v(x, t)$  is an invariant surface of (6), i.e., it corresponds to the invariant surface condition  $\tau v_t + \xi(x, t, v)v_x = \eta(x, t, v)$ , then  $u = v(x, t)$  is an invariant solution of Eq. (1) relating to the infinitesimal generator of (6).

For the symmetry of  $V_1$  of (18), we give the characteristic equation

$$\frac{dx}{x} = \frac{\alpha dt}{4t} = \frac{du}{-u},\tag{19}$$

and the corresponding invariants are

$$z = xt^{-\frac{\alpha}{4}}, \quad u = t^{-\frac{\alpha}{4}} f(z).\tag{20}$$

**Theorem 2** The time fractional nonlinear fourth-order evolution equation is reduced to the nonlinear ordinary differential equation(ODE) of time fractional order in the following by means of similarity transformation  $u = t^{-\frac{\alpha}{4}} f(z)$  and the similarity variable  $z = xt^{-\frac{\alpha}{4}}$ ,

$$\left(P_{\frac{\alpha}{4}}^{1-\frac{5\alpha}{4}, \alpha} f\right)(z) + h_1 f_z f_{zz} + h_2 f^2 f_{zz} + h_3 f(f_z)^2 + h_4 f_{zzzz} = 0,\tag{21}$$

with the Erdélyi-Kober fractional differential operator  $P_{\delta}^{\tau, \alpha}$  of order<sup>[34]</sup>

$$(P_{\delta}^{\tau, \alpha} g)(z) := \prod_{j=0}^{m-1} \left( \tau + j - \frac{1}{\delta} z \frac{d}{dz} \right) (K_{\delta}^{\tau + \alpha, m - \alpha} g)(z), \quad z > 0, \quad \delta > 0, \quad \alpha > 0,\tag{22}$$

$$m = \begin{cases} [\alpha] + 1, & \alpha \notin \mathbb{N}, \\ \alpha, & \alpha \in \mathbb{N}, \end{cases}\tag{23}$$

where

$$(K_{\delta}^{\tau, \alpha} g)(z) := \begin{cases} \frac{1}{\Gamma(\alpha)} \int_1^{\infty} (v-1)^{\alpha-1} v^{-(\tau+\alpha)} g(zv^{\frac{1}{\delta}}) dv, & \alpha > 0, \\ g(z), & \alpha = 0, \end{cases}\tag{24}$$

is the Erdélyi-Kober fractional integral operator.

**Proof** Let  $n - 1 < \alpha < n$ ,  $n = 1, 2, \dots$ , then we obtain the similarity transformation of Riemann-Liouville fractional derivatives as follows

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^n}{\partial t^n} \left[ \frac{1}{\Gamma(n - \alpha)} \int_0^t (t - s)^{n - \alpha - 1} s^{-\frac{\alpha}{4}} f(xs^{-\frac{\alpha}{4}}) ds \right]. \quad (25)$$

If  $v = \frac{t}{s}$ , we get

$$\begin{aligned} \frac{\partial^\alpha u}{\partial t^\alpha} &= \frac{\partial^n}{\partial t^n} \left[ t^{n - \frac{5\alpha}{4}} \frac{1}{\Gamma(n - \alpha)} \int_1^\infty (v - 1)^{n - \alpha - 1} v^{-(n + 1 - \frac{5\alpha}{4})} f(zv^{\frac{\alpha}{4}}) dv \right] \\ &= \frac{\partial^n}{\partial t^n} \left[ t^{n - \frac{5\alpha}{4}} \left( K_{\frac{\alpha}{4}}^{1 - \frac{\alpha}{4}, n - \alpha} f \right) (z) \right]. \end{aligned} \quad (26)$$

Considering the relation of  $z = xt^{-\frac{\alpha}{4}}$ , we get

$$t \frac{\partial}{\partial t} \Phi(z) = tx \left( -\frac{\alpha}{4} \right) t^{-\frac{\alpha}{4} - 1} \Phi'(z) = -\frac{\alpha}{4} z \frac{d}{dz} \Phi(z). \quad (27)$$

Hence, the simplification of the above equation is

$$\begin{aligned} \frac{\partial^n}{\partial t^n} \left[ t^{n - \frac{5\alpha}{4}} \left( K_{\frac{\alpha}{4}}^{1 - \frac{\alpha}{4}, n - \alpha} f \right) (z) \right] &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[ \frac{\partial}{\partial t} \left( t^{n - \frac{5\alpha}{4}} \left( K_{\frac{\alpha}{4}}^{1 - \frac{\alpha}{4}, n - \alpha} f \right) (z) \right) \right] \\ &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[ t^{n - \frac{5\alpha}{4}} \left( n - \frac{5\alpha}{4} - \frac{\alpha}{4} z \frac{d}{dz} \left( K_{\frac{\alpha}{4}}^{1 - \frac{\alpha}{4}, n - \alpha} f \right) (z) \right) \right]. \end{aligned} \quad (28)$$

Continuously, we have

$$\begin{aligned} \frac{\partial^n}{\partial t^n} \left[ t^{n - \frac{5\alpha}{4}} \left( K_{\frac{\alpha}{4}}^{1 - \frac{\alpha}{4}, n - \alpha} f \right) (z) \right] &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[ \frac{\partial}{\partial t} \left( t^{n - \frac{5\alpha}{4}} \left( K_{\frac{\alpha}{4}}^{1 - \frac{\alpha}{4}, n - \alpha} f \right) (z) \right) \right] \\ &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[ t^{n - \frac{5\alpha}{4}} \left( n - \frac{5\alpha}{4} - \frac{\alpha}{4} z \frac{d}{dz} \left( K_{\frac{\alpha}{4}}^{1 - \frac{\alpha}{4}, n - \alpha} f \right) (z) \right) \right] \\ &= \dots = t^{-\frac{5\alpha}{4}} \prod_{j=0}^{n-1} \left( 1 - \frac{5\alpha}{4} + j - \frac{\alpha}{4} z \frac{d}{dz} \right) \left( K_{\frac{\alpha}{4}}^{1 - \frac{\alpha}{4}, n - \alpha} f \right) (z). \end{aligned} \quad (29)$$

Considering the fractional differential operator of Erdélyi-Kober fractional operator of (22), we get

$$\frac{\partial^n}{\partial t^n} \left[ t^{n - \frac{5\alpha}{4}} \left( K_{\frac{\alpha}{4}}^{1 - \frac{\alpha}{4}, n - \alpha} f \right) (z) \right] = t^{-\frac{5\alpha}{4}} \left( P_{\frac{\alpha}{4}}^{1 - \frac{5\alpha}{4}, \alpha} f \right) (z). \quad (30)$$

Thus,

$$\frac{\partial^\alpha u}{\partial t^\alpha} = t^{-\frac{5\alpha}{4}} \left( P_{\frac{\alpha}{4}}^{1 - \frac{5\alpha}{4}, \alpha} f \right) (z). \quad (31)$$

Therefore, it's fascinating that Eq. (1) reduces to the fractional-order ODE

$$\left( P_{\frac{\alpha}{4}}^{1 - \frac{5\alpha}{4}, \alpha} f \right) (z) + h_1 f_z f_{zz} + h_2 f^2 f_{zz} + h_3 f(f_z)^2 + h_4 f_{zzzz} = 0. \quad (32)$$

## 4 Conservation laws

### 4.1 Preliminaries of conservation laws

According to the Riemann-Liouville left-sided time-fractional derivative<sup>[35–37]</sup>, we have

$${}_0D_t^\alpha u = D_t^n ({}_0I_t^{n-\alpha} u), \tag{33}$$

in Eq. (1), where  $D_t$  is the operator concerning  $t$  of differentiation,  $n = [\alpha] + 1$ . In addition, the definition of  ${}_0I_t^{n-\alpha} u$  is given as follows.

**Definition 3**<sup>[22]</sup> The left-sided time-fractional integral of order  ${}_0I_t^{n-\alpha} u$  is defined by

$$({}_0I_t^{n-\alpha} u)(x, t) = \frac{1}{\Gamma(n - \alpha)} \int_0^t \frac{u(\theta, x)}{(t - \theta)^{1-n+\alpha}} d\theta, \tag{34}$$

where,  $\Gamma(z)$  is the Gamma function.

**Definition 4** A conserved vector is said to obey the conservation law, if it satisfies the following conservation equation

$$D_t(C^t) + D_x(C^x) = 0, \tag{35}$$

where  $C^t = C^t(t, x, u, \dots)$ ,  $C^x = C^x(t, x, u, \dots)$ .

Firstly, we suppose that Eq. (1) has the following formal Lagrangian

$$L = v(x, t) [u_t^\alpha + h_1 u_x u_{xx} + h_2 u^2 u_{xx} + h_3 u u_x^2 + h_4 u_{xxxx}], \tag{36}$$

where,  $v(x, t)$  is a new dependent variable. In view of the previous Lagrangian, we get an action integral as follows

$$\int_0^T \int_\Omega L(x, t, u, v, D_t^\alpha(u), u_x, \dots) dx dt. \tag{37}$$

Similar to the case of integer-order nonlinear differential equations<sup>[30–31]</sup>, the adjoint equation is available. Therefore, we get the conclusion that the Euler-Lagrange equation of nonlinear fourth-order evolution equation Eq. (1) with adjoint equation in the following

$$\frac{\delta L}{\delta u} = 0, \tag{38}$$

and the definition of the Euler-Lagrange operator is obtained as below.

**Definition 5**<sup>[30–31]</sup> The Euler-Lagrange operator is defined by

$$\frac{\delta}{\delta u} = \frac{\partial}{\partial u} + (D_t^\alpha)^* \frac{\partial}{\partial D_t^\alpha u} - D_x \frac{\partial}{\partial u_x} + D_x^2 \frac{\partial}{\partial u_{xx}} - D_x^3 \frac{\partial}{\partial u_{3x}} + D_x^4 \frac{\partial}{\partial u_{xxxx}}, \tag{39}$$

where  $(D_t^\alpha)^*$  is the adjoint operator of  $(D_t^\alpha)$ .

For the Riemann-Liouville differential operators

$$(D_t^\alpha)^* = (-1)^n I_T^{n-\alpha} (D_t^n) \equiv {}_t^C D_T^\alpha, \tag{40}$$

here,  $I_T^{n-\alpha}$  is the right-sided operator of fractional integration of order  $n - \alpha$ , and  $I_T^{n-\alpha}$  is defined by

$$I_T^{n-\alpha} f(t, x) = \frac{1}{\Gamma(n - \alpha)} \int_t^T \frac{f(\tau, x)}{(\tau - t)^{1+\alpha-n}} d\tau, \quad n = [\alpha] + 1. \tag{41}$$

We find that Eq. (1.1) with the Riemann-Liouville fractional derivative satisfies the conservation equation of Eq. (33) as follows

$$C^t = D_t^{n-1}({}_0I_t^{n-\alpha}u), \quad C^x = h_4u_{xxx} + h_2u^2u_x + \frac{h_1}{2}u_x^2 \quad (42)$$

when,  $h_3 = 2h_2$ .

In case of two independent variables  $t, x$ , and one dependent variable  $u(x, t)$ , we get

$$\bar{X} + D_t(\tau)l + D_x(\xi)l = W \frac{\delta}{\delta u} + D_t N^t + D_x N^x, \quad (43)$$

where  $l$  is the identity operator,  $\frac{\delta}{\delta u}$  shows the Euler-Lagrange operator, and the Noether operators are provided by  $N^t, N^x$  respectively. In addition,  $\bar{X}$  is defined by

$$\bar{X} = \tau \frac{\partial}{\partial t} + \xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial u} + \xi_\alpha^0 \frac{\partial}{\partial D_t^\alpha u} + \eta^x \frac{\partial}{\partial u_x} + \eta^{xx} \frac{\partial}{\partial u_{xx}} + \eta^{xxx} \frac{\partial}{\partial u_{xxx}} + \eta^{xxxx} \frac{\partial}{\partial u_{xxxx}}, \quad (44)$$

and

$$W = \eta - \tau u_t - \xi u_x. \quad (45)$$

Applying the Riemann-Liouville time-fractional derivative to Eq. (1), we obtain the following operator  $N^t$  defined in [30–31]

$$N^t = \tau l + \sum_{k=0}^{n-1} (-1)^k {}_0D_t^{\alpha-1-k}(W) D_t^k \frac{\partial}{\partial ({}_0D_t^\alpha u)} - (-1)^n J \left( W, D_t^n \frac{\partial}{\partial ({}_0D_t^\alpha u)} \right), \quad (46)$$

where  $J$  is the integral,

$$J(f, g) = \frac{1}{\Gamma(n-\alpha)} \int_0^t \int_t^T \frac{f(\tau, x)g(\mu, x)}{(\mu-\tau)^{\alpha+1-n}} d\mu dt. \quad (47)$$

The operator  $N^x$  is defined by

$$\begin{aligned} N^x = & \xi l + W \left( \frac{\partial}{\partial u_x} - D_x \frac{\partial}{\partial u_{xx}} + D_x^2 \frac{\partial}{\partial u_{xxx}} - D_x^3 \frac{\partial}{\partial u_{xxxx}} \right) \\ & + D_x(W) \left( \frac{\partial}{\partial u_{xx}} - D_x \frac{\partial}{\partial u_{xxx}} + D_x^2 \frac{\partial}{\partial u_{xxxx}} \right) \\ & + D_x^2(W) \left( \frac{\partial}{\partial u_{xxx}} - D_x \frac{\partial}{\partial u_{xxxx}} \right) \\ & + D_x^3 \frac{\partial}{\partial u_{xxxx}}. \end{aligned} \quad (48)$$

No matter what the solutions and generator  $X$  of Eq. (1) are, we obtain

$$(\bar{X}L + D_t(\tau)L + D_x(\xi)L)|_{(1)} = 0. \quad (49)$$

It is easy to find that Euler-Lagrange (38) of Eq. (1) is equal to zero. Hence, the conservation law resulted from the right-hand side of the equality is given by<sup>[30]</sup>

$$D_t(N^t L) + D_x(N^x L) = 0. \quad (50)$$



Comparing (35) and (50), we find that the following components of conserved vectors is always valid obviously

$$C^t = N^t L, \quad C^x = N^x L. \tag{51}$$

**4.2 Conservation laws of time fractional foam drainage equation**

In this section, the conservation laws of Eq. (1) is available.

**Theorem 3** The components of conserved vectors about fourth-order evolution equation is given by the following two cases

**Case 1** when  $\alpha \in (0, 1)$

$$\begin{aligned} V_1 : C_1^t &= {}_0D_t^{\alpha-1}(W_1)v + J(W_1, v_t) \\ &= {}_0D_t^{\alpha-1}\left(-u - \frac{4t}{\alpha}u_t - xu_x\right)v + J\left(-u - \frac{4t}{\alpha}u_t - xu_x, v_t\right), \\ C_1^x &= -\left(u + \frac{4t}{\alpha}u_t + xu_x\right)[2v(h_3 - h_2)uu_x - v_x(h_1u_x + h_2u^2)] \\ &\quad - v\left(2u_x + \frac{4t}{\alpha}u_{xr} \right) \end{aligned} \tag{52}$$



where the functions of  $W_i$ ,  $i = 1, 2$  have the form

$$W_1 = -u - \frac{4t}{\alpha}u_t - xu_x, \quad W_2 = -u_x. \quad (56)$$

**Proof** Substituting (46) and (48) into Eq. (1), we get the conserved vector as below.

For the case of  $\alpha \in (0, 1)$ ,

$$\begin{aligned} C_i^t &= \tau L + (-1)^0 {}_0D_t^{\alpha-1}(W_i)D_t^0 \frac{\partial L}{\partial ({}_0D_t^\alpha u)} - (-1)J \left( W_i, D_t \frac{\partial L}{\partial ({}_0D_t^\alpha u)} \right) \\ &= {}_0D_t^{\alpha-1}(W_i)v + J(W_i, v_t), \end{aligned} \quad (57)$$

and

$$\begin{aligned} N^x &= \xi l + W \left( \frac{\partial}{\partial u_x} - D_x \frac{\partial}{\partial u_{xx}} + D_x^2 \frac{\partial}{\partial u_{xxx}} - D_x^3 \frac{\partial}{\partial u_{xxxx}} \right) \\ &\quad + D_x(W) \left( \frac{\partial}{\partial u_{xx}} - D_x \frac{\partial}{\partial u_{xxx}} + D_x^2 \frac{\partial}{\partial u_{xxxx}} \right) \\ &\quad + D_x^2(W) \left( \frac{\partial}{\partial u_{xxx}} - D_x \frac{\partial}{\partial u_{xxxx}} \right) \\ &\quad + D_x^3 \frac{\partial}{\partial u_{xxxx}}. \end{aligned} \quad (58)$$

For the case of  $\alpha \in (1, 2)$ ,

$$\begin{aligned} C_i^t &= \tau L + (-1)^0 {}_0D_t^{\alpha-1}(W_i)D_t^0 \frac{\partial L}{\partial ({}_0D_t^\alpha u)} - (-1)^1 J \left( W_i, D_t^1 \frac{\partial L}{\partial ({}_0D_t^\alpha u)} \right) \\ &\quad + (-1)^1 {}_0D_t^{\alpha-2}(W_i)D_t^1 \frac{\partial L}{\partial ({}_0D_t^\alpha u)} - (-1)^2 J \left( W_i, D_t^2 \frac{\partial L}{\partial ({}_0D_t^\alpha u)} \right) \\ &= {}_0D_t^{\alpha-1}(W_i)v + J(W_i, v_t) - {}_0D_t^{\alpha-2}(W_i)v_t - J(W_i, v_{tt}), \end{aligned} \quad (59)$$

and

$$\begin{aligned} N^x &= \xi l + W \left( \frac{\partial}{\partial u_x} - D_x \frac{\partial}{\partial u_{xx}} + D_x^2 \frac{\partial}{\partial u_{xxx}} - D_x^3 \frac{\partial}{\partial u_{xxxx}} \right) \\ &\quad + D_x(W) \left( \frac{\partial}{\partial u_{xx}} - D_x \frac{\partial}{\partial u_{xxx}} + D_x^2 \frac{\partial}{\partial u_{xxxx}} \right) \\ &\quad + D_x^2(W) \left( \frac{\partial}{\partial u_{xxx}} - D_x \frac{\partial}{\partial u_{xxxx}} \right) \\ &\quad + D_x^3 \frac{\partial}{\partial u_{xxxx}}, \end{aligned} \quad (60)$$

here

$$D_x = \frac{\partial}{\partial x} + u_x \frac{\partial}{\partial u} + v_x \frac{\partial}{\partial v} + u_{xx} \frac{\partial}{\partial u_x} + u_{xt} \frac{\partial}{\partial u_t} + v_{xx} \frac{\partial}{\partial v_x} + v_{xt} \frac{\partial}{\partial v_t} + \dots \quad (61)$$

and  $u = u(x, t)$ ,  $v = v(x, t)$ .

For the operators  $V_1 = x \frac{\partial}{\partial x} + \frac{4t}{\alpha} \frac{\partial}{\partial t} - u \frac{\partial}{\partial u}$ ,  $V_2 = \frac{\partial}{\partial x}$ , referring to (45), we get the Lie characteristics functions,  $W_1 = -u - \frac{4t}{\alpha}u_t - xu_x$ ,  $W_2 = -u_x$ .

In accordance with the fractional generalizations of the Riemann-Liouville differential operators<sup>[30]</sup>, we have the specific form of the time fractional fourth-order evolution equation as follows. When  $\alpha \in (0, 1)$ , we substitute (36) along with (56) into (57) and (58). When  $\alpha \in (1, 2)$ , we substitute (36) and (56) into (59) and (60), and see that Theorem 3 is 1.00,0.00,0.00. The proof is completed.

## 5 Conclusions and discussions

In this paper, we study the time fractional nonlinear fourth-order evolution equation by way of Lie symmetry. Based on known results, we conclude the corresponding vector fields. In addition, these corresponding vector fields are further applied in order to construct the symmetry reductions of  $\frac{\partial^\alpha u}{\partial t^\alpha} + h_1 u_x u_{xx} + h_2 u^2 u_{xx} + h_3 u u_x^2 + h_4 u_{xxxx}, 0 < \alpha \leq 2$ . In the end, we obtain the conservation laws. However, we still have a lot of problems to research, such as how can we study the time-space fractional equation? How to solve the fractional differential equations with more independent and dependent variables? These are our future work.

## References:

- [ 1 ] Bluman G W, Anco S. Symmetry and integration methods for differential equations [M]. New York: Springer-Verlag, 2002.
- [ 2 ] Ibragimov N H. CRC handbook of Lie group analysis of differential equations: Symmetries, exact solutions and conservation laws [M]. New York: CRC Press, 1994.
- [ 3 ] Lakshmanan M, Kaliappan P. Lie transformations, nonlinear evolution equations, and Painlevé forms [J]. Journal of mathematical physics, 1983, 24(4):795–806.
- [ 4 ] Olver P J. Applications of Lie groups to differential equations [M]. New York: Springer-Verlag, 1986.
- [ 5 ] Ovsiannikov L V. Group analysis of differential equations [M]. Ames W F, Transl. New York: Academic Press, 1982.
- [ 6 ] Tian S F. The mixed coupled nonlinear Schrödinger equation on the half-line via the Fokas method [J]. Proceedings of the Royal Society A: Mathematical Physical and Engineering Sciences, 2016, 472:20160588.
- [ 7 ] Wang D S, Yin Y B. Symmetry analysis and reductions of the two-dimensional generalized Benney system via geometric approach [J]. Computers & Mathematics with Applications 2016, 71:748-757.
- [ 8 ] Wang D S, Wei X Q. Integrability and exact solutions of a two-component Korteweg-de Vries system [J]. Applied Mathematics Letters 2016, 51:60-67.
- [ 9 ] Tian S F, Zhang H Q. On the integrability of a generalized variable-coefficient Kadomtsev-Petviashvili equation [J]. Journal of Physics A: Mathematical and Theoretical 2012, 45(5):055203(29pp).
- [10] Tian S F, Zhang Y F, Feng B L, et al. On the Lie algebras, generalized symmetries and darboux transformations of the fifth-order evolution equations in shallow water [J]. Chinese Annals of Mathematics, Series B, 2015, 36(4):543-560.
- [11] Tian S F. Initial-boundary value problems for the general coupled nonlinear Schrödinger equation on the interval via the Fokas method [J]. Journal of Differential Equations 2017, 262(1):506-558.
- [12] Tian S F, Zhang H Q. On the integrability of a generalized variable-coefficient forced Korteweg-de Vries equation in fluids [J]. Studies in Applied Mathematics 2014, 132(3):212-246.
- [13] Diethelm K. The analysis of fractional differential equations [M]. New Nork: Springer dordrecht heidelberg, 2010.
- [14] Liang Y J, Chen W. A survey on computing Lévy stable distributions and a new matlab toolbox [J]. Signal Processing, 2013, 93(1):242-251.
- [15] Hu S, Chen W, Gou X F. Modal analysis of fractional derivative damping model of frequency-dependent viscoelastic soft matter [J]. Advances in Vibration Engineering, 2011, 10:187-196.
- [16] El-Sayed A M A, Gaber M. The Adomian decomposition method for solving partial differential equations of fractal order in finite domains [J]. Physics Letters A, 2006, 359(3):175-182.
- [17] Gazizov R K, Kasatkin A A, Lukashchuk S Y. Continuous transformation groups of fractional differential equations [J]. Vestnik USATU, 2007, 9(3):21.
- [18] Gazizov R K, Kasatkin A A, Lukashchuk S Y. Symmetry properties of fractional diffusion equations [J]. Physica Scripta, 2009(T136):014016.
- [19] Sahadevan R, Bakkyaraj T. Invariant analysis of time fractional generalized Burgers and Korteweg-de Vries equations [J]. Journal of mathematical analysis and applications, 2012, 393(2):341-343.

- [20] Wang G W, Xu T Z. Invariant analysis and exact solutions of nonlinear time fractional Sharma-Tasso-Olver equation by Lie group analysis [J]. *Nonlinear Dynamics*, 2014, 76:571-580.
- [21] Huang Q, Zhdanov R. Symmetries and exact solutions of the time fractional Harry-Dym equation with Riemann-Liouville derivative [J]. *Physic A*, 2014, 409:110-118.
- [22] Lukashchuk S Y, Makunin A V. Group classification of nonlinear time-fractional diffusion equation with a source term [J]. *Applied Mathematics and Computation*, 2015, 257:335-343.
- [23] Noether E. Invariant variationa problems [J]. *Transport Theory and Statistical Physics*, 1971, 1(3):183-207.
- [24] Frederico G S F, Torres D F M. A formulation of Noethers theorem for fractional problems of the calculus of variations [J]. *Journal of Mathematical Analysis and Applications*. 2007, 334(2):834-846.
- [25] Atanackovic T M, Konjik S, Pilipovic S, et al. Variational problems with fractional derivatives: invariance conditions and Nöthers theorem [J]. *Nonlinear Analysis*, 2009, (5-6):1504-1517.
- [26] Malinowska A B. A formulation of the fractional Noether-type theorem for multidimensional Lagrangians [J]. *Applied Mathematics Letters*, 2012, 25(11):1941-1946.
- [27] Odziejewicz T, Malinowska A B, Torres D F M. Noethers theorem for fractional variational problems of variable order [J]. *Central European Journal of Physics*, 2013, 11(6):691-701.
- [28] Bourdin L, Cresson J, Greff I. A continuous/discrete fractional Noethers theorem [J]. *Communications in Nonlinear Science and Numerical Simulation*, 2013, 18(4):878-887.
- [29] Ibragimov N H. A new conservation theorem [J]. *Journal of Mathematical Analysis and Applications*, 2007, 333(1):311-328.
- [30] Lukashchuk S Y. Conservation laws for time-fractional subdiffusion and diffusion-wave equations [J]. *Nonlinear Dynamics*, 2015, 80(1):791-802.
- [31] Gazizov R K, Ibragimov N H, Lukashchuk S Y. Nonlinear self-adjointness, conservation laws and exact solutions of time-fractional Kompaneets equations [J]. *Communications in Nonlinear Science and Numerical Simulation*, 2015, 23(1-3):153-163.
- [32] Nikolai A K, Maria V D. Polygons of differential equations for finding exact solutions [J]. *Chaos, Solitons and Fractals*, 2007, 33(5):1480-1496.
- [33] Ibragimov N H. Transformation groups applied to mathematical physics [M]. New York: Springer-Verlag, 1985.
- [34] Kiryakova V S. Generalised fractional calculus and applications [M]. New York: CRC Press, 1993.
- [35] Baleanu D, Diethelm K, Scalas E, et al. Fractional calculus: Models and numerical methods [M]. Singapore: World Scientific Publishing Company, 2012.
- [36] Samko S, Kilbas A, Marichev O. Fractional integrals and derivatives: Theory and applications [M]. London: Gordon and Breach, 1993.
- [37] Kilbas A A A, Srivastava H M, Trujillo J J. Theory and applications of fractional differential equations [M]. Amsterdam: Elsevier Science Limited, 2006.

(责任编辑: 顾浩然)